

The Solar Structure and the Five-minute Oscillation

R. Scuflaire, M. Gabriel, and A. Noels

Institut d'Astrophysique, Université de Liège, Avenue de Cointe 5, B-4200 Ougrée-Liège, Belgium

Received September 29, accepted December 10, 1980

Summary. A sequence of solar envelope models with convective zones of variable thickness has been computed. The corresponding theoretical eigenfrequencies of radial and nonradial modes were then compared with recent observations. This favours a model with a convective zone containing 7.5% of the mass and extending over 34% of the radius. Such a convective zone is much deeper than in the standard solar model.

Key words: solar oscillations

1. Introduction

The five-minute solar oscillation has been the object of detailed studies.

The observations of Deubner (1975), Rhodes et al. (1977), Deubner et al. (1979) have simultaneously resolved the motion of the solar surface in frequency and in horizontal wave-number. They have brought out the f -mode and the first few p -modes of large spherical harmonics degree ℓ (typically a few hundreds). Modes of this type have been computed for solar models by Ulrich (1970), Ando and Osaki (1975, 1977), Ulrich and Rhodes (1977), Lubow et al. (1980), Berthomieu et al. (1980), and Rhodes and Ulrich (1980). They have found that the theoretical frequencies are sensitive mainly to the depth of the convective zone and that these observations are better accounted for with a convective zone deeper than was previously thought.

Claverie et al. (1979, 1980) succeeded in observing oscillations near five minutes in integral sunlight. They interpreted them as low ℓ , high overtones pulsations. Christensen-Dalsgaard and Gough (1980) showed that they must be modes with ℓ values lower than 4.

We have computed a sequence of solar envelope models. They are described in Sect. 2. The eigenfrequencies are computed and compared with those observed by Claverie et al. (1980) for low ℓ values and with those of Deubner et al. (1979) for high ℓ values. The consequences of these comparisons for the solar structure are then discussed.

2. The Models

A sequence of evolution has been computed with the Henyey code (Henyey et al., 1964, 1965) for a solar model with an initial composition given by $X = 0.7417$, $Z = 0.02$. The equation of state is

given by Bodenheimer et al. (1965). The nuclear reaction rates are those of Fowler et al. (1967). For the opacity coefficient, we took the values of Cox and Stewart (1970), modified at low temperatures according to Alexander (1975) to take into account the effect of molecules. The model corresponding to the solar age ($4.5 \cdot 10^9$ yr), model 1, has the correct (solar) radius ($R_{\odot} = 6.96 \cdot 10^{10}$ cm) and luminosity ($L_{\odot} = 3.82 \cdot 10^{33}$ erg s $^{-1}$) when a value of 1.85 is chosen for ℓ/H , the ratio of the mixing-length to the pressure scale height. We were particularly careful in the computation of the external layers and the model includes a model of chromosphere whose temperature profile is given by Vernazza et al. (1973). The model of chromosphere extends up to the discontinuity of temperature located at 2400 km above the photosphere. We assumed that the pressure vanishes at this point. From the bottom of the convective zone up to this point the model includes about two hundred points. As most evolution codes do, the model has been broken into two parts for the computation: an interior and an envelope. The mesh size is smaller in the envelope than in the interior. At the transition between these two parts (which corresponds to a temperature of about $3 \cdot 10^5$ K) we required a relative precision of 0.001 on the fit.

The other models considered here (models 2 to 5) are solar envelope models comprising 80% of the mass. They are obtained by an inward integration alone, they do not fit an interior model. The temperature at their basis is of the order of 10^7 K. They all have the solar radius and luminosity and the same chemical composition as our model 1. They differ by the thickness of the convective zone, corresponding to different values of ℓ/H . We shall not discuss here the difficult problem of which type of interior can match these envelope models. Table 1 gives some

Table 1. Some characteristics of the models. ℓ/H is the ratio of the mixing-length in the convective zone to the pressure scale height, r , ρ , and T the radius, the density and the temperature at the bottom of the convective zone, m_c the mass of the convective zone and τ the life-time of Li^7 in the convective zone

Model no.	ℓ/H	r (R_{\odot})	m_c (M_{\odot})	ρ (g cm $^{-3}$)	T (10^6 K)	τ (10^9 yr)
1	1.85	0.744	0.014	0.106	1.88	91,200
2	2.4	0.704	0.031	0.239	2.30	421
3	3	0.674	0.053	0.396	2.63	16.3
4	4	0.649	0.084	0.613	2.92	1.36
5	5	0.633	0.107	0.781	3.10	0.342
X	3.7	0.66	0.075	0.548	2.83	2.74

Send offprint requests to: R. Scuflaire

Table 2. The frequencies ν (mHz) of modes of order n (p_n) of low ℓ values are given for model 1 in the range 2.3 to 4 mHz

n	$\ell=0$	$\ell=1$	$\ell=2$	$\ell=3$
15			2.3365	2.3888
16	2.3524	2.4139	2.4706	2.5231
17	2.4865	2.5480	2.6052	2.6584
18	2.6213	2.6832	2.7407	2.7941
19	2.7566	2.8187	2.8766	2.9302
20	2.8924	2.9545	3.0122	3.0663
21	3.0280	3.0904	3.1485	3.2028
22	3.1640	3.2266	3.2850	3.3398
23	3.3004	3.3634	3.4220	3.4771
24	3.4372	3.5003	3.5595	3.6149
25	3.5744	3.6378	3.6969	3.7528
26	3.7116	3.7754	3.8349	3.8908
27	3.8492	3.9131	3.9728	
28	3.9869			

Table 3. The frequencies ν (mHz) of modes of order n (p_n) of low ℓ values are given for model 5 in the range 2.3 to 4 mHz

n	$\ell=0$	$\ell=1$	$\ell=2$	$\ell=3$
15			2.3486	2.4056
16	2.3552	2.4205	2.4825	2.5395
17	2.4899	2.5534	2.6162	2.6735
18	2.6245	2.6866	2.7499	2.8073
19	2.7591	2.8198	2.8833	2.9414
20	2.8935	2.9535	3.0167	3.0756
21	3.0277	3.0878	3.1502	3.2099
22	3.1619	3.2224	3.2842	3.3446
23	3.2962	3.3576	3.4183	3.4794
24	3.4302	3.4932	3.5530	3.6143
25	3.5645	3.6289	3.6882	3.7495
26	3.6991	3.7649	3.8239	3.8849
27	3.8339	3.9010	3.9600	
28	3.9692			

characteristics for the five models of the sequence. The life-time of lithium in the convective zone has been computed according to Fowler et al. (1975). It varies approximately as T^{-24} (T being the temperature at the bottom of the convective zone) and the ratio of the life-time to its local value at the bottom of the convective zone is approximately 14.1 for all models. Our values are in good agreement with those of Ulrich and Rhodes (1977).

3. Modes of Low ℓ Values

Modes of low ℓ values were computed as described in Boury et al. (1975). In order to have good eigenvalues and eigenfunctions we used a fourth order integration method with a variable stepsize independent of the mesh size of the model. The stepsize is chosen as the integration is proceeding in such a way to keep the discretization error small. Halving the stepsize, we have obtained, in a few test cases, an estimation of the error due to the discretization process. In our computation the relative error on the frequency amounts to a few 10^{-6} . This result is achieved with about one thousand points. This error is quite negligible com-

Table 4. For all the models and for each value of ℓ the table gives the frequency ν (mHz) closest to 3 mHz and the mean separation $\Delta\nu$ (μ Hz) between frequencies in the range 2.3 to 4 mHz

Model no.	$\ell=0$	$\ell=1$	$\ell=2$	$\ell=3$
1	3.0280	2.9545	3.0122	2.9302
	136.21	136.29	136.36	136.55
2	3.0326	2.9612	3.0205	2.9405
	135.88	135.71	135.91	136.10
3	3.0325	2.9608	3.0220	2.9433
	135.50	135.20	135.34	135.58
4	3.0300	2.9569	3.0198	2.9430
	134.91	134.75	134.64	134.90
5	3.0277	2.9535	3.0167	2.9414
	134.50	134.59	134.28	134.48

pared to the error arising from the lack of an interior model (for models 2 to 5), a problem we discuss just below.

As models 2 to 5 are envelope models we met a difficulty in the application of the internal boundary condition. Applying a rigid boundary condition at the bottom of these models can lead to rather large relative errors on the frequencies which may amount to 4 or 5%, as was shown by a test on model 1. Then we replaced the missing interior by an artificial distribution of density given by

$$\rho(r) = a + br^2,$$

where the parameters a and b are determined so that ρ and m be continuous. (The continuity of the pressure automatically results from the inward integration of the hydrostatic equilibrium equation.) We have verified on model 1 that this process introduces relative errors on the frequencies at most of the order of $2 \cdot 10^{-4}$.

Christensen-Dalsgaard and Gough (1980) have shown that the discrete frequencies observed by Claverie et al. (1979, 1980) in the range 2.3 to 4 mHz correspond to modes with $\ell < 4$ and fall into nearly degenerate groups satisfying

$$\nu_{\ell,n} \simeq \nu_{\ell+2,n-1},$$

where ℓ is the degree of the spherical harmonics and n the order of the overtone. The approximate relation

$$\nu_{\ell+1,n} \simeq \frac{1}{2}(\nu_{\ell,n} + \nu_{\ell,n+1})$$

also holds for these modes. These relations are confirmed by our computations. Tables 2 and 3 give the computed frequencies for our extreme models. The twenty-five frequencies observed by Claverie et al. (1980) may be alternately identified with a mode of even ℓ ($n=16$ to 28 for $\ell=0$ or 15 to 27 for $\ell=2$) and a mode of odd ℓ ($n=16$ to 27 for $\ell=1$ or 15 to 26 for $\ell=3$). Therefore the observed frequencies fall into two spectra.

As the frequencies of these two spectra as well as the computed spectrum for a given ℓ are nearly equally spaced, each spectrum may be represented by two parameters, the value of one frequency ν (we have chosen the frequency closest to 3 mHz) and the mean separation $\Delta\nu$ between frequencies in the range from 2.3 to 4 mHz. Table 4 gives these parameters for all models. In Fig. 1 we have represented the two observed spectra with their uncertainties by the crosses and each continuous line represents the computed spectra of a given ℓ for the sequence of models. We notice that the mean separation between frequencies is rather sensitive to the




Fig. 1. ν and $\Delta\nu$ are respectively the frequency closest to 3 mHz and the mean separation between frequencies. The two crosses represent the observed spectra, respectively of odd and even values of ℓ . The continuous curves represent for each value of ℓ the computed spectra for the sequence of models. For one of these curves the numbers label the models

thickness of the convective zone. Modes of $\ell = 1$ and $\ell = 2$ would match satisfactorily the observations for a model whose characteristics are given under the entry *X* in Table 1.

4. Modes of High ℓ Values

Modes of high ℓ values (typically a few hundreds) were computed as described by Gabriel and Noels (1976), however the variables were chosen as in Gabriel and Scuflaire (1979). As δr varies as $r^{\ell-1}$ in the central regions, the effect of the missing inner regions on the eigenfrequencies is completely negligible.

Solar oscillation modes of high ℓ values have been computed by a number of authors (Ulrich, 1970; Ando and Osaki, 1977; Lubow et al., 1980; Berthomieu et al., 1980). The comparisons with observations (Rhodes et al., 1977; Berthomieu et al., 1980; Rhodes and Ulrich, 1980) rule out a shallow convective zone: it must at least extend down to $0.7R_{\odot}$.

Our own computations (Tables 5 and 6) confirm the necessity of a deep convective zone. Figure 2 shows that our model with the deepest convective zone (model 5) agree better with the observations of Deubner et al. (1979) than our standard solar model (model 1).

5. Conclusion

As the observations of the five-minute oscillation are becoming more refined, the requirements imposed on the models of the external layers of the sun are getting more severe. We have shown that they can be interpreted with models having a normal solar composition, it is not necessary to resort to models with a low *Z* in the interior as first suggested by Claverie et al. (1979) from the low ℓ observations. Both the low and high ℓ observations require a model with a deep convective envelope extending down to $0.66R_{\odot}$ and containing 7.5% of the mass. This result is compatible with the Li^7 abundance observed in the sun as its lifetime remains greater than 10^9 yr.

Table 5. The frequencies ν (mHz) of modes of order n of high ℓ values are given for model 1. The asterisk indicates the chromospheric mode

n	$\ell=200$	$\ell=400$	$\ell=600$	$\ell=800$
0	1.4272	2.0077	2.4540	2.8304
1	1.9563	2.5672	3.0653	3.5192
2	2.3895	3.1082	3.6918	3.8880*
3	2.7662	3.6489	3.8016*	4.2360
4	3.1368	3.7264*	4.3189	4.7562
5	3.4877	4.1611	4.6296	4.9366

Table 6. The frequencies ν (mHz) of modes of order n of high ℓ values are given for model 5. The asterisk indicates the chromospheric mode

n	$\ell=200$	$\ell=400$	$\ell=600$	$\ell=800$
0	1.4268	2.0086	2.4546	2.8302
1	1.9572	2.5733	3.0527	3.4939
2	2.3907	3.0893	3.6452	3.8869*
3	2.7655	3.5932	3.7815*	4.1512
4	3.1180	3.6820*	4.2305	4.5597
5	3.4558	4.0804	4.4251	4.8061




Fig. 2. The power contour in this ℓ, ν diagram is taken from Deubner et al. (1979). The crosses indicate the computed modes of model 1. When they differ significantly from those of model 1 the frequencies of model 5 have been represented by triangles. The chromospheric mode has been omitted

The problem we have now to face is that the standard solar model requires a much thinner convective zone than found here. It is tempting to conclude that after the neutrinos observations, the solar oscillations show that something is wrong in the standard solar models.

Acknowledgements. We are indebted to A. Claverie, G. R. Isaak, C. P. McLeod, and H. B. van der Raay and T. Roca Cortes for sending us their observations before publication. It is a pleasure to thank R. K. Ulrich, who referred this paper, for his advice.

References

- Alexander, D.R.: 1975, *Astrophys. J. Suppl.* **29**, 363
 Ando, H., Osaki, Y.: 1975, *Publ. Astron. Soc. Japan* **27**, 581
 Ando, H., Osaki, Y.: 1977, *Publ. Astron. Soc. Japan* **29**, 221
 Berthomieu, G., Copper, A.J., Gough, D.O., Osaki, Y., Provost, J., Rocca, A.: 1980, in *Nonradial and nonlinear stellar pulsation*, eds. H. A. Hill and W. A. Dziembowski, Springer, Berlin, Heidelberg, New York, p. 307
 Bodenheimer, P., Forbes, J.E., Gould, N.L., Henyey, L.G.: 1965, *Astrophys. J.* **141**, 1019
 Boury, A., Gabriel, M., Noels, A., Scuflaire, R., Ledoux, P.: 1975, *Astron. Astrophys.* **41**, 279
 Christensen-Dalsgaard, J., Gough, D.O.: 1980, *Monthly Notices Roy. Astron. Soc.* (in press)
 Claverie, A., Isaak, G.R., McLeod, C.P., Van der Raay, H.B.: 1979, *Nature* **282**, 591
 Claverie, A., Isaak, G.R., McLeod, C. P., Van der Raay, H.B., Roca Cortes, T.: 1980, *Astron. Astrophys.* **91**, L9
 Cox, A.N., Stewart, J.N.: 1970, *Astrophys. J. Suppl.* **19**, 243
 Deubner, F.L.: 1975, *Astron. Astrophys.* **44**, 371
 Deubner, F.L., Ulrich, R.K., Rhodes, E.J.: 1979, *Astron. Astrophys.* **72**, 177
 Fowler, W.A., Caughlan, G.R., Zimmerman, B.A.: 1967, *Ann. Rev. Astron. Astrophys.* **5**, 525
 Fowler, W.A., Caughlan, G.R., Zimmerman, B. A.: 1975, *Ann. Rev. Astron. Astrophys.* **13**, 69
 Gabriel, M., Noels, A.: 1976, *Astron. Astrophys.* **53**, 149
 Gabriel, M., Scuflaire, R.: 1979, *Acta Astron.* **29**, 135
 Henyey, L.G., Forbes, J.E., Gould, N.L.: 1964, *Astrophys. J.* **139**, 306
 Henyey, L.G., Vardya, M.S., Bodenheimer, P.: 1965, *Astrophys. J.* **142**, 841
 Lubow, S.H., Rhodes, E.J., Ulrich, R.K.: 1980, in *Nonradial and nonlinear stellar pulsation*, eds. H.A. Hill, and W. A. Dziembowski, Springer, Berlin, Heidelberg, New York, p. 300
 Rhodes, E.J., Ulrich, R.K., Simon, G.W.: 1977, *Astrophys. J.* **218**, 901
 Rhodes, E.J., Ulrich, R.K.: 1980, *Bull. Amer. Astron. Soc.* **12**, 475
 Ulrich, R.K.: 1970, *Astrophys. J.* **162**, 993
 Ulrich, R.K., Rhodes, E.J.: 1977, *Astrophys. J.* **218**, 521
 Vernazza, J.E., Avrett, E.H., Loeser, R.: 1973, *Astrophys. J.* **184**, 605