



Solar-Like Oscillations in a Massive Star

Kévin Belkacem, *et al.*
Science **324**, 1540 (2009);
DOI: 10.1126/science.1171913

The following resources related to this article are available online at www.sciencemag.org (this information is current as of June 19, 2009):

Updated information and services, including high-resolution figures, can be found in the online version of this article at:

<http://www.sciencemag.org/cgi/content/full/324/5934/1540>

Supporting Online Material can be found at:

<http://www.sciencemag.org/cgi/content/full/324/5934/1540/DC1>

This article **cites 8 articles**, 1 of which can be accessed for free:

<http://www.sciencemag.org/cgi/content/full/324/5934/1540#otherarticles>

This article appears in the following **subject collections**:

Astronomy

<http://www.sciencemag.org/cgi/collection/astronomy>

Information about obtaining **reprints** of this article or about obtaining **permission to reproduce this article** in whole or in part can be found at:

<http://www.sciencemag.org/about/permissions.dtl>

4. P. V. Nguyen, T. Abel, E. R. Kandel, *Science* **265**, 1104 (1994).
5. K. C. Martin *et al.*, *Cell* **91**, 927 (1997).
6. U. Frey, R. G. Morris, *Nature* **385**, 533 (1997).
7. A. Casadio *et al.*, *Cell* **99**, 221 (1999).
8. K. C. Martin, *Curr. Opin. Neurobiol.* **14**, 305 (2004).
9. M. A. Sutton, E. M. Schuman, *Cell* **127**, 49 (2006).
10. A. Govindarajan, R. J. Kelleher, S. Tonegawa, *Nat. Rev. Neurosci.* **7**, 575 (2006).
11. O. Steward, W. B. Levy, *J. Neurosci.* **2**, 284 (1982).
12. L. E. Ostroff, J. C. Fiala, B. Allwardt, K. M. Harris, *Neuron* **35**, 535 (2002).
13. S. J. Tang *et al.*, *Proc. Natl. Acad. Sci. U.S.A.* **99**, 467 (2002).
14. J. Eberwine, B. Belt, J. E. Kacharmina, K. Miyashiro, *Neurochem. Res.* **27**, 1065 (2002).
15. R. Moccia *et al.*, *J. Neurosci.* **23**, 9409 (2003).
16. J. Zhong, T. Zhang, L. M. Bloch, *BMC Neurosci.* **7**, 17 (2006).
17. M. M. Poon, S. H. Choi, C. A. Jamieson, D. H. Geschwind, K. C. Martin, *J. Neurosci.* **26**, 13390 (2006).
18. T. Suzuki, Q. B. Tian, J. Kurumitsu, T. Kawai, S. Endo, *Neurosci. Res.* **57**, 61 (2007).
19. H. Kang, E. M. Schuman, *Science* **273**, 1402 (1996).
20. K. M. Huber, J. C. Roder, M. F. Bear, *J. Neurophysiol.* **86**, 321 (2001).
21. G. Aakalu, W. B. Smith, N. Nguyen, C. Jiang, E. M. Schuman, *Neuron* **30**, 489 (2001).
22. C. Job, J. Eberwine, *Nat. Rev. Neurosci.* **2**, 889 (2001).
23. W. Ju *et al.*, *Nat. Neurosci.* **7**, 244 (2004).
24. K. F. Raab-Graham, P. C. Haddick, Y. N. Jan, L. Y. Jan, *Science* **314**, 144 (2006).
25. V. F. Castellucci, S. Schacher, *Prog. Brain Res.* **86**, 105 (1990).
26. S. L. Mackey *et al.*, *Proc. Natl. Acad. Sci. U.S.A.* **84**, 8730 (1987).
27. V. Lyles, Y. Zhao, K. C. Martin, *Neuron* **49**, 349 (2006).
28. D. L. Glanzman, E. R. Kandel, S. Schacher, *Neuron* **3**, 441 (1989).
29. J. F. Brunet, E. Shapiro, S. A. Foster, E. R. Kandel, Y. Iino, *Science* **252**, 856 (1991).
30. J. Y. Hu, Y. Chen, S. Schacher, *J. Neurosci.* **27**, 11712 (2007).
31. J. Y. Hu, F. Wu, S. Schacher, *J. Neurosci.* **26**, 1026 (2006).
32. N. G. Gurskaya *et al.*, *Nat. Biotechnol.* **24**, 461 (2006).
33. Materials and methods are available as supporting material on *Science Online*.
34. K. Liu, J. Y. Hu, D. Wang, S. Schacher, *J. Neurobiol.* **56**, 275 (2003).
35. Z. Guan *et al.*, *Cell* **111**, 483 (2002).
36. L. Santarelli, P. Montarolo, S. Schacher, *J. Neurobiol.* **31**, 297 (1996).
37. D. Cai, S. Chen, D. L. Glanzman, *Curr. Biol.* **18**, 920 (2008).
38. We thank S. Braslow and K. Cadenas for assistance with image analysis; R. Grambo for assistance with figures; D. Black, C. Heusner, E. Meer, and L. Zipursky for critical reading of the manuscript; and G. Weinmaster and Martin laboratory members for helpful discussions. This work was supported by NIH grant NS045324, a W. M. Keck Foundation Young Scholar Award and Eleanor Leslie Term Chair from the UCLA Brain Research Institute (to K.C.M.), Canadian Institute of Health Research grant MT-15121 (to W.S.S.), and a fellowship from the Nakajima Foundation (to S.K.M.).

Supporting Online Material

www.sciencemag.org/cgi/content/full/1173205/DC1
Materials and Methods
Figs. S1 to S17
References

9 March 2009; accepted 5 May 2009
Published online 14 May 2009;
10.1126/science.1173205
Include this information when citing this paper.

REPORTS

Solar-Like Oscillations in a Massive Star


Kévin Belkacem,^{1,2*} Réza Samadi,¹ Marie-Jo Goupil,¹ Laure Lefèvre,¹ Frédéric Baudin,³ Sébastien Deheuvels,¹ Marc-Antoine Dupret,^{1,2} Thierry Appourchaux,³ Richard Scuflaire,² Michel Auvergne,¹ Claude Catala,¹ Eric Michel,¹ Andrea Miglio,² Josefina Montalban,² Anne Thoul,² Suzanne Talon,⁴ Annie Baglin,¹ Arlette Noels²

Seismology of stars provides insight into the physical mechanisms taking place in their interior, with modes of oscillation probing different layers. Low-amplitude acoustic oscillations excited by turbulent convection were detected four decades ago in the Sun and more recently in low-mass main-sequence stars. Using data gathered by the Convection Rotation and Planetary Transits mission, we report here on the detection of solar-like oscillations in a massive star, V1449 Aql, which is a known large-amplitude (β Cephei) pulsator.

Stars burn hydrogen into helium through nuclear fusion during most of their life. Once the central hydrogen gets exhausted, the helium core starts contracting, and hydrogen-shell burning takes over as the main energy source. The subsequent evolution depends mostly on a star's mass at birth but also on the physical mechanisms occurring during the hydrogen-burning phase. For instance, transport of chemical elements determines the helium core size, which is crucial for the evolution of stars. Transport pro-

cesses such as turbulence and those induced by rotation are not fully understood and are still poorly modeled, but stellar seismology can pro-

Fig. 1. Fourier spectrum of prewhitened light curve obtained from the quasi-uninterrupted 150 days of observations, with a duty cycle of 90%, of the star V1449 Aql by CoRoT, showing structures that are reproduced over the 100- to 250- μ Hz interval. (Inset) Enlarged part of the spectrum showing a typical solar-like structure. Below 100 μ Hz, we enter the bulk regime of unstable modes (fig. S1), and the possible existence of many such modes



in this frequency domain then makes the deciphering of unstable versus stable modes quite delicate. Hence, to remain conservative we restrict the discussion to frequencies above 100 μ Hz.

¹Laboratoire d'Études Spatiales et d'Instrumentation en Astrophysique, CNRS (UMR 8109), Observatoire de Paris, Place J. Janssen, F-92195 Meudon, France; associated with Université Pierre et Marie Curie and Université Denis Diderot. ²Institut d'Astrophysique et de Géophysique de l'Université de Liège, Allée du 6 Août 17-B 4000 Liège, Belgium. ³Institut d'Astrophysique Spatiale, Université Paris-Sud 11 and CNRS (UMR 8617), Batiment 121, F-91405 Orsay, France. ⁴Réseau Québécois de Calcul de Haute Performance, Université de Montréal, Casier Postal 6128, Succursale Centre-ville, Montréal, Québec H3C 3J8, Canada.

*To whom correspondence should be addressed. E-mail: kevin.belkacem@ulg.ac.be

iron-opacity bump in such a massive star induces the existence of a convective zone in the upper layers (4), which could be responsible for the excitation of the detected modes.

Our results are based on the quasi-uninterrupted (meaning, a duty cycle of 90%) light curve obtained over 150 days with the Convection Rotation and Planetary Transits (CoRoT) (5–7) Centre National d'Etudes Spatiales (CNES) space mission (Fig. 1). The dominant opacity-driven mode is located at 63.5 μHz with an amplitude of 3.9×10^4 parts per million (ppm). We looked for stochastically excited modes with frequencies above 100 μHz ; below this limit, the existence of several opacity-driven modes makes the analysis more difficult. No signal is found

above 250 μHz . In the frequency range 100 to 250 μHz , there are broad structures with a width of several μHz , with low amplitudes of hundreds of parts per million and well above the noise level, which is around 1 ppm.

These structures are not the result of instrumental effects [supporting online material (SOM text)]. To verify that they are not related to the existence of the opacity-driven modes, we carried out prewhitening (SOM text), which suppresses the influence of the dominant peaks and their harmonics in the relevant frequency domain as well as related aliases because of the observational interruptions. We validated our prewhitening method through numerical simulations (SOM text).





Fig. 2. Time-frequency diagram, using a Morlet wavelet with a 20-day width (8). **(Top)** Solar-like mode in the prewhitened light curve shown in the inset of Fig. 1, which exhibits a time-dependent behavior and a spreading over several μHz . **(Bottom)** For comparison, the second harmonic of the dominant peak, associated with the opacity-driven mode in the unprewhitened light curve.

Fig. 3. Autocorrelation of the power-density spectrum associated with V1449 Aql. The red curve corresponds to the autocorrelation smoothed with a boxcar average width of 0.60 μHz . The autocorrelation has been computed between 130 and 300 μHz . The domain below 130 μHz is not considered to avoid the low-order modes, which in general depart from the regular spacing expected with the high-order p modes.



The modes associated with the broad structures have a finite lifetime, in contrast with opacity-driven modes, which are coherent oscillations and therefore appear as sinus cardinal functions in the Fourier spectrum. The amplitudes of these oscillations vary stochastically in time, again in contrast with the stationary property of the dominant opacity-driven mode and its harmonics. Their power is intermittent in time and dispersed in terms of frequency (Fig. 2). Such behavior, typical of solar-like modes (8), confirms the stochastic nature of the structures. In contrast, the temporal behavior of the second harmonic of the fundamental opacity-driven mode, which lies in the same frequency interval, is centered on a single frequency (Fig. 2, bottom). The widths of the detected high-frequency modes show that they are damped, which is a signature of modes excited by turbulent convection.

We looked for regularly spaced patterns in the Fourier spectrum, which are a characteristic signature of those modes. An autocorrelation of the Fourier spectrum shows periodicities centered around 5, 14, and 27 μHz (Fig. 3), indicating the existence of periodicities in the power spectrum.

Theoretical calculations show that these properties, interpreted as damped acoustic modes excited by turbulent convection, are compatible with solar-like oscillations of a massive main sequence star. We carried out numerical simulations using a 10-solar mass stellar model that is appropriate for V1449 Aql in that it corresponds to the observational constraints obtained from ground-based observations (9). A comparison between the theoretical and observational autocorrelations shows that the observed frequency spectrum is compatible with the presence of modes of angular degrees $l=0, 1, \text{ and } 2$, characterized by a large frequency separation around 27 μHz , with 1- μHz widths and a rotational splitting of 2.5 μHz related to a rotation with an axis inclined by 90° with respect to the line of sight (SOM text).

Mode amplitudes obtained from theoretical computations of the line width (10) and the energy supplied in the mode by turbulent convection (11) reach several tens of parts per million, which is well above the CoRoT detection threshold and in agreement with observations. Our calculations show that excitation by the turbulent convective motions associated with the iron-opacity bump in the upper layers of the star is efficient. This driving is operative when the convective time scale of energy-bearing eddies is close to the modal period, which explains why modes in the frequency range of 100 to 250 μHz are observed.

In summary, we showed that the broad structures at high frequencies detected in the CoRoT Fourier spectrum of the star V1449 Aql are not the result of instrumental effects, are independent of the opacity-driven modes, and present regularly spaced patterns that are characteristic of high-frequency acoustic modes. These structures have the theoretically expected properties of solar-like oscillations: modes excited by turbulent convection.

References and Notes

1. C. Waelkens *et al.*, *Astron. Astrophys.* **330**, 215 (1998).
2. K. Uytterhoeven *et al.*, *J. Phys. Conf. Ser.* **118**, 012077 (2008).
3. W. A. Dziembowski, A. A. Pamiatnykh, *Mon. Not. R. Astron. Soc.* **262**, 204 (1993).
4. M. Cantiello *et al.*, *Astron. Astrophys.* **499**, 279 (2009).
5. The CoRoT space mission, launched on 27 December 2006, has been developed and is operated by CNES, with the contribution of Austria, Belgium, Brazil, the European Space Agency (ESA) (Research and Scientific Support Department and Science Programme), Germany, and Spain.
6. M. Auvergne *et al.*, *Astron. Astrophys.*, <http://arXiv.org/abs/0901.2206>.
7. E. Michel *et al.*, *Science* **322**, 558 (2008).
8. F. Baudin, A. Gabriel, D. Gibert, *Astron. Astrophys.* **285L**, 29 (1994).
9. T. Morel, C. Aerts, *CoAst* **150**, 201 (2007).
10. M. A. Dupret, *Astron. Astrophys.* **366**, 166 (2001).
11. K. Belkacem *et al.*, *Astron. Astrophys.* **478**, 163 (2008).
12. K.B. acknowledges financial support from Liège University through the Subside Fédéral pour la Recherche. Part of this research was funded by the

Belgian Prodex-ESA. L.L. has been partly supported by CNES. A.T. is Chercheur Qualifié at the Fonds National de la Recherche Scientifique (FNRS) and A.M. is Chargé de recherches FNRS.

Supporting Online Material

www.sciencemag.org/cgi/content/full/324/5934/1540/DC1

SOM Text
Figs. S1 to S5
References

5 February 2009; accepted 13 May 2009
10.1126/science.1171913

Colloidal Quantum-Dot Photodetectors Exploiting Multiexciton Generation

Vlad Sukhovatkin, Sean Hinds, Lukasz Brzozowski, Edward H. Sargent*

Multiexciton generation (MEG) has been indirectly observed in colloidal quantum dots, both in solution and the solid state, but has not yet been shown to enhance photocurrent in an optoelectronic device. Here, we report a class of solution-processed photoconductive detectors, sensitive in the ultraviolet, visible, and the infrared, in which the internal gain is dramatically enhanced for photon energies E_{photon} greater than 2.7 times the quantum-confined bandgap E_{bandgap} . Three thin-film devices with different quantum-confined bandgaps (set by the size of their constituent lead sulfide nanoparticles) show enhancement determined by the bandgap-normalized photon energy, $E_{\text{photon}}/E_{\text{bandgap}}$, which is a clear signature of MEG. The findings point to a valuable role for MEG in enhancing the photocurrent in a solid-state optoelectronic device. We compare the conditions on carrier excitation, recombination, and transport for photoconductive versus photovoltaic devices to benefit from MEG.

Multiexciton generation (MEG) refers to the creation of two or more electron-hole pairs per absorbed photon in a semiconductor (1). Colloidal quantum-dot materials in which MEG has been reported experimentally include PbS and PbSe (2), PbTe (3), CdSe (4), and Si (5). In bulk semiconductors, carrier multiplication has been observed repeatedly over the past five decades, both in elemental semiconductors such as germanium (6) and silicon (7) and also in lead chalcogenides (8), including the infrared-bandgap bulk semiconductor PbS (9). In the past year, experiments that carefully account for processes such as photoionization of nanoparticles during spectroscopic studies have evidenced the production of more than one exciton per photon (10) in colloidal quantum dots, with yields ranging from 1.1 to 2.4 excitons per photon (10) when the photon energy exceeds the MEG threshold near $\sim E_{\text{photon}}/E_{\text{bandgap}} > 2.7$ (11), where E_{photon} is the photon energy and E_{bandgap} is the quantum-confined bandgap.

MEG has been reported, based on all-optical spectroscopic data, not only in solution but also in thin solid films; however, in spite of numerous attempts with materials systems and photon energies reported to manifest MEG, neither the external quantum efficiency (EQE) nor the internal quantum efficiency (IQE) of the photocurrent in a device has been shown to exceed 100% (12–20).

In particular, one careful and systematic study (21) recently explored whether a key signature of MEG—an IQE of greater than unity—was observable in the photocurrent of a low-bandgap PbSe colloidal quantum-dot photovoltaic device. Once reflection and absorption were carefully taken into account, IQEs approaching, but not exceeding, 100% were reported.

Recent reports (22) suggest that in some of the earlier spectroscopic studies, the apparent quantum yield of MEG was enhanced by photoionization in the presence of multiple excitons. The sequence of steps is depicted in Fig. 1. The generation of two excitons within one quantum dot (Fig. 1B) produces efficient Auger recombination; one exci-

ton recombines, and one carrier associated with the other exciton is excited high within its band (Fig. 1C). Photoionization, in this instance known as Auger-assisted ionization (AAI), may occur when this excited charge carrier becomes trapped at or near the quantum dot's surface, resulting in a nanoparticle that possesses a long-lived net charge (Fig. 1D). The energetic Auger electron has a higher probability of being captured to a trap than does an already-thermalized electron (23) because it can more easily surmount the energetic barrier (such as a thin oxide on the nanoparticle surface), restricting access to a surface trap state.

Subsequent photogeneration of even a single exciton then results in the presence of a trion (an exciton plus a charge) that recombines rapidly, masquerading as MEG in recombination dynamics-based studies. Thus, MEG's time-resolved spectroscopic signatures are enhanced by photoionization. It should be emphasized that, at the low intensities of interest in MEG investigations, photoionization alone cannot masquerade as MEG and serves only to amplify the apparent quantum yield if MEG is already present.

Unfortunately, MEG combined with photoionization provides no advantages over MEG alone in the harvest of photovoltaic energy. Indeed, because trions resulting from photoionization accelerate recombination, they render even more challenging the extraction of MEG photocurrent from a photovoltaic device. To observe the benefits of MEG in the current extracted from a photovoltaic device, charge separation must occur before




Fig. 1. MEG accompanied by photoionization. (A) Bands and trap levels for the quantum dots that were used. E_c is the quantum-confined conduction band edge, E_t is the trap energy, $E_{g_{qd}}$ is the quantum-confined bandgap, and $E_{g_{bulk}}$ is the constituent semiconductor's bulk bandgap. (B) Generation of a pair of excitons via photon absorption followed by carrier multiplication. (C) Auger-induced excitation of an electron to a higher-lying level concomitant with recombination of the other exciton. (D) Efficient trapping of the excited electron.

Department of Electrical and Computer Engineering, University of Toronto, Toronto, ON M5S 3G4, Canada.

*To whom correspondence should be addressed. E-mail: ted.sargent@utoronto.ca



www.sciencemag.org/cgi/content/full/324/5934/1540/DC1

Supporting Online Material for

Solar-like Oscillations in a Massive Star

Kévin Belkacem, Réza Samadi, Marie-Jo Goupil, Laure Lefèvre, Frédéric Baudin, Sébastien Deheuvels, Marc-Antoine Dupret, Thierry Appourchaux, Richard Scuflaire, Michel Auvergne, Claude Catala, Eric Michel, Andrea Miglio, Josefina Montalban, Anne Thoul, Suzanne Talon, Annie Baglin, Arlette Noels

Published 19 June 2009, *Science* **324**, 1540 (2009)

DOI: 10.1126/science.1171913

This PDF file includes:

SOM Text

Figs. S1 to S5

References

Supporting online material

Solar-like oscillations in a massive star

Kévin Belkacem^{1,2}, Réza Samadi¹, Marie-Jo Goupil¹, Laure Lefèvre¹, Frédéric Baudin³, Sébastien Deheuvels¹, Marc-Antoine Dupret^{1,2}, Thierry Appourchaux³, Richard Scuflaire², Michel Auvergne¹, Claude Catala¹, Eric Michel¹, Andrea Miglio², Josefina Montalbán², Anne Thoul², Suzanne Talon⁴, Annie Baglin¹, Arlette Noels²

¹LESIA, CNRS (UMR 8109), Observatoire de Paris, pl. J. Janssen, F-92195 Meudon, France, associé Université Pierre et Marie Curie, Université Denis Diderot

²Institut d'Astrophysique et de Géophysique de l'Université de Liège, Allée du 6 Août 17 – B 4000 Liège, Belgium

³Institut d'Astrophysique Spatiale (IAS), Université Paris-Sud 11 and CNRS (UMR 8617), Batiment 121, F-91405 Orsay, France

⁴Réseau québécois de calcul de haute performance (RQCHP), Université de Montréal, C.P. 6128, succursale centre-ville, Montréal, Québec, H3C 3J8

I) Data analysis

a) Instrumental effects

As a first test to verify that the structures seen in the frequency domain [100; 250] μHz of the Fourier spectrum of V1449 Aql (HD180642) are not due to some instrumental perturbations, we have considered the fainter (apparent visual magnitude $m_v=9.14$ for HD181072, and $m_v=8.27$ for V1449 Aql (SI)) CoRoT target HD181072, which is in the same observation run and same CCD.

HD181072 does not exhibit structures comparable to those observed in V1449 Aql in the abovementioned frequency region. Its frequency spectrum only shows a residual of the orbital frequency with an amplitude around 9.8 ppm and a flat/white noise with an amplitude of 0.6 ppm in 150 days, far below the amplitudes of the structures we are interested in. Furthermore, neither the star background nor the satellite jitter show the structures observed in the stellar V1449 Aql.

b) Prewhitening procedure

To remove the influence of the dominant peaks and associated perturbation due to observational interruptions, we performed a prewhitening of the light curve (hereafter LC) obtained by CoRoT (Fig SI). The highest peaks in the time-series are fitted and then removed from the time series. Within each frequency range of 2 μHz , peaks are selected whenever their amplitudes are at least five times the local mean of the power spectrum. A fit to the time series is performed, based on a Levenberg-Marquardt method with a least-square minimization (S2, S3, S4). A total of 91 peaks were withdrawn. We find that the large structures in the power spectrum, which we identify as solar-like modes, are unaltered by the prewhitening procedure; their existence is therefore independent of the low frequency opacity-driven modes.

c) Validation using numerical simulations

In order to validate our prewhitening procedure as well as to theoretically investigate the influence of the combined effects of the unstable dominant modes and the observational interruptions, we built two simulated LC using our CoRoT light-curve simulator (*S5*). The first simulation is a temporal signal built by including only the low-frequency unstable modes that have been detected in the LC of V1449 Aql down to the amplitude of 460 ppm. In the second simulation, we also included a set of solar-like oscillations. The frequencies and line widths of these solar-like oscillations are computed using the non-adiabatic numerical code MAD (*S6*) and a stellar model- built with the evolutionary numerical code CLES (*S7*) - that matches the effective temperature and the luminosity of the star (see section II for details). Amplitudes of the included solar-like modes are set to a level representative of the high-frequency stable modes detected in V1449 Aql, that is 300 ppm. Each simulated LC is sampled with the same duty-cycle than that associated with the observations of V1449 Aql and noise is added to the resulting time series. Finally, we filled the gaps in the same way as done by the CoRoT data treatment chain (*S8*).

The prewhitened Fourier spectrum associated with the first simulated LC does not show any structures in the frequency domain we are interested in. This demonstrates that the combined effects of the unstable dominant modes, the window and the filling procedure do not produce spurious structures that could erroneously be interpreted as solar-like oscillations.

The second LC, which includes solar-like modes, has been prewhitened to remove the unstable modes in the same manner than done for V1449 Aql. The Fourier spectrum associated with the second simulated LC is presented in Fig. S2. The simulated solar-like oscillations are recovered in the Fourier spectrum associated with the second simulated LC. This demonstrates that the clean procedure leaves existing solar-like oscillations unaltered.

II) Modelling of solar-like modes

a) Auto-correlation of HD180642

Solar like oscillations are acoustic oscillations whose frequencies are expected to follow more or less closely the asymptotic regime (*S9*). In order to search for such a signature of regular p-mode spacing, we computed the autocorrelation of the Fourier spectrum in the frequency domain 130 μHz – 300 μHz . The domain below 130 μHz is excluded so as to avoid the low-frequency (low-radial order) modes, which in general significantly depart from the regular spacing expected for the high frequency p modes. The result is shown in Fig. S3, which clearly reveals three patterns, one centred on $\nu_0 = 5 \mu\text{Hz}$, the second around $\nu_1 = 13.5 \mu\text{Hz}$ and the last one around $\nu_2 = 27 \mu\text{Hz}$. The second pattern is centered at a frequency equal to half ν_2 , that is $\nu_1 = \nu_2/2$. The autocorrelation function then shows that a regular frequency spacing exists in the Fourier spectrum of the light-curve of V1449 Aql. This is the expected signature of asymptotic p modes with an associated large separation of either $\nu_2 = 27 \mu\text{Hz}$ or half its value, ν_1 . We also see that the peak at ν_2 in the autocorrelation function has several components that are distant from ν_2 by a multiple of 2.5 μHz . The same feature,

although less pronounced, is also observed in the pattern centred on ν_1 . The presence of the peak at $\nu_0=5 \mu\text{Hz}$ can be interpreted as the signature of mode splitting induced by the rotation of the star.

b) Simulated auto-correlation

An important issue is to verify that the properties of high frequency p modes, which are derived from the observed spectrum, are -at least roughly- consistent with theoretical expectations for solar-like p modes in a massive star. We therefore carried out a modelling of V1449 Aql, which is not fully optimized but sufficient for the present purpose. Future works will be dedicated to an optimized modelling of V1449 Aql.

To simulate an autocorrelation that can be compared with the one derived from the observed V1449 Aql LC, we computed a model that approximately matches the effective temperature of the star (*S1*) ($T_{\text{eff}}=24500 \pm 1000 \text{ K}$, $\log g = 3.45 \pm 0.15$). A ten-solar-mass model is obtained, with an effective temperature of 24075 K and a gravity of $\log g=3.88$, using the CLES evolutionary code (*S7*). The associated p-mode frequencies present a large separation of $27 \mu\text{Hz}$, as suggested by the observed autocorrelation (Fig. S3). In building the simulated light curve, we included modes with angular degree $l=0, 1, 2$ and 3 . We assumed the same theoretical value, $\sigma=1 \mu\text{Hz}$, obtained with the code MAD (*S6*) for all line-widths of the modes. The radial modes were all given amplitude of 300 ppm, which is representative of the structures observed for V1449 Aql. The amplitude of the non-radial modes were set up to a 300 ppm value modulated by mode visibility factors. We further assumed a constant rotational splitting σ and defined i as the inclination angle between the rotation axis and the line of sight. From our set of theoretical frequencies, we simulate a LC using the CoRoT light-curve simulator¹⁵ and computed the associated power density spectrum and the autocorrelation function.

We investigated different values of σ and i . An agreement with the observations is found for a rotational splitting of $\sigma= 2.5 \mu\text{Hz}$ and an inclination angle of $i=90^\circ$. The result is shown in Fig S4. The pattern at ν_0 is reproduced. Our theoretical result shows that including p-modes split by a rotation of $2.5 \mu\text{Hz}$ with an inclination angle of $i=90^\circ$ gives an autocorrelation that is compatible with the observed one. In order for the reader to estimate the quality of the agreement, we also show two additional autocorrelation functions in Fig. S5. In the first case we assume $i=45^\circ$ and a splitting of $2.5 \mu\text{Hz}$ (top panel), while in the other case we assume $i=90^\circ$ with no splitting, i.e. keeping only the central component of the rotationally split multiplets (bottom panel). These autocorrelation functions do not reproduce the observed structures seen in Fig. S3.

These results provide a theoretical support to our assumption that the structures observed on the autocorrelation function of V1449 Aql are compatible with the presence of a regular p-mode spectrum. We also find that the main features of the observed autocorrelation function can be reproduced by assuming an internal rotation of $2.5 \mu\text{Hz}$ and a rotation axis that has an inclination of $i\sim 90^\circ$ with respect to the observed direction. However, a definite determination of the values of angle i and the rotation frequency is dedicated to a forthcoming work together with an optimized modeling.




Fig S1: Fourier spectrum of prewhitened light curve obtained from the quasi-uninterrupted 150 days of observations, with a duty-cycle of 90%, of the star V1449 Aql by CoRoT, showing structures that are reproduced. Below 100 μHz , we enter the bulk regime of unstable modes and the possible existence of many such modes in this frequency domain then makes the deciphering of unstable versus stable modes quite delicate. Below 50 μHz , one notes an increase of the power toward low frequencies that is not yet identified as due to coloured noise, and/or g modes, and/or rotation. (Inset) Enlarged part of the spectrum.




Fig S2: Power spectral density of the *simulated* LC that includes both unstable modes and solar-like oscillations. In black: prewhitened LC. In red: original LC




Fig S3: Autocorrelation of the power density spectrum associated with the observed lightcurve of V1449 Aql. The autocorrelation has been computed between 130 μHz and 300 μHz .




Fig S4: Autocorrelation function of the *simulated* power density spectrum associated with the stellar model that matches the effective temperature of V1449 Aql and that has a large separation of 27 μHz . It includes the modes $l=0,1$ and 2, with $i=90^\circ$ and a splitting of 2.5 μHz .




Fig S5: Autocorrelation function of simulated power density spectrum associated with the stellar model that matches the effective temperature of V1449 Aql and that has a large separation of 27 μHz . At the top panel, the simulation includes the modes $l=0,1$ and 2, with $i=45^\circ$ and a splitting of 2.5 μHz . At the bottom panel, the simulation includes the modes $l=0,1$ and 2, with $i=90^\circ$ and no splitting.

REFERENCES AND NOTES

- S1. Morel, T., Aerts, C., An abundance study of the B-type targets for the asteroseismology programme of the CoRoT mission, *CoAst*, **150**, 201 (2007)
- S2. Lenz, P., Breger, M., Period04 User Guide, *CoAst*, **146**, 53-136 (2005)
- S3. Koen, C., The analysis of indexed astronomical time series - V. Fitting sinusoids to high-speed photometry, *MNRAS*, **309**, 769-802 (1999)
- S4. Martinez, P., Koen, C., Period Searching by Least Squares in Sparsely Sampled Lightcurves with Non-Sinusoidal Oscillations, *MNRAS*, **267**, 1039 (1994)
- S5. Baudin, F., Samadi, R., Appourchaux, T., Michel, E., Simu-LC : a Light-Curve simulator for CoRoT, Contribution to "The Corot book" published by ESA (SP 1306; Eds: M. Fridlund, A. Baglin, L. Conroy and J. Lochard), 2007
- S6. Dupret, M.A., Nonradial nonadiabatic stellar pulsations: A numerical method and its application to a beta Cephei model, *Astron.and Astrophys.*, **366**, 166-173 (2001)
- S7. Scuflaire, R. et al., CLÉS, Code Liégeois d'Évolution Stellaire, *Astrophysics and Space Science*, **316**, 83-91 (2008)
- S8. Samadi, R. et al., "The Corot Book: Chap. V.5/ Extraction of the photometric information: corrections", Contribution to "The Corot book" published by ESA (SP 1306; Eds: M. Fridlund, A. Baglin, L. Conroy and J. Lochard), 2007
- S9. Christensen-Dalsgaard, J., Helioseismology, *Reviews of Modern Physics*, **74**, 1073-1129 (2002)